§ 301.38-2

States Department of Agriculture,² and that are listed as rust-resistant under § 301.38-2 (a)(2) and (a)(3).

Rust-susceptible plants. All plants of the genera Berberis, Mahoberberis, and Mahonia species not listed as rust-resistant under §301.38-2 (a)(2) and (a)(3).

Regulated article. Any article listed in $\S 301.38-2$ (a)(1) through (a)(4) of this subpart or otherwise designated as a regulated article in accordance with $\S 301.38-2$ (a)(5) of this subpart.

Seedling. Any plant of the genera Berberis, Mahoberberis, and Mahonia grown from seed and having less than 2 years' growth.

State. The District of Columbia, Puerto Rico, the Northern Mariana Islands, or any State, territory or possession of the United States.

Two years' growth. The growth of a plant during all growing seasons of 2 successive calendar years.

[54 FR 32791, Aug. 10, 1989; 54 FR 38494, Sept. 18, 1989; 67 FR 8178, Feb. 22, 2002]

§301.38-2 Regulated articles.

- (a) The following are regulated articles: 3
- (1) All seedlings and plants of less than 2 years' growth of the genus *Berberis*.
- (2) All plants, seeds, fruits, and other plant parts capable of propagation from the following rust-resistant *Berberis* species.

```
B. aggregata x B. wilsoniae 'Pirate
King'
```

B. 'Amstelveen'

B. aridocalida

B. beaniana

B. buxifolia

B. buxifolia nana

B. calliantha

B. candidula

B. candidula 'Amstelveen'

B. candidula×B. verruculosa

'Amstelveen' B. cavallieri

D. cavaillell

B. chenaulti

B. chanaulti 'Apricot Queen'

B. circumserrata

B. concinna

B. coxii

B. darwini

B. dasystachya

B. dubia

B. feddeana

B. formosana

B. franchetiana

B. gagnepainii

B. gagnepaini 'Chenault'

B. gilgiana

B. gladwynensis

B. gladwynensis 'William Penn'

B. gyalaica

B. heterophylla

B. horvathi

B. hybrido-gagnepaini

B. insignis

B. integerrima 'Wallichs Purple'

B. julianae

B. julianae 'Nana'

B. julianae 'Spring Glory'

B. koreana

B. koreana×B. thunbergii hybrid Bailsel

B. koreana×B. thunbergii hybrid Tara

B. lempergiana

B. lepidifolia

B. linearifolia

B. linearifolia var. 'Orange King'

B. lologensis

B. lologensis 'Mystery Fire'

B. manipurana

B. media 'Park Jewel'

B. media 'Red Jewel'

B. mentorensis

B. pallens

B. poirettii 'BJG 073', 'MTA'

B. potanini

B. Renton

B. replicata

B. sanguinea B. sargentiana

²Testing is performed by the Agricultural Research Service of USDA as follows: In a greenhouse, the suspect plant, or test subject, is placed under a screen with a control plant, i.e., a known rust-susceptible variety of Berberis, Mahoberberis, or Mahonia. Infected wheat stems, a primary host of black stem rust, are placed on top of the screen. The plants are moistened and maintained in 100% humidity, causing the spores to swell and fall on the plants lying under the screen. The plants are then observed for 7 days at 20-80% relative humidity. This test procedure is repeated 12 times. If in all 12 tests, the rustsusceptible plant shows signs of infection after 7 days and the test plants do not, USDA will declare the test plant variety rust-resistant. The tests must be performed on new growth, just as the leaves are unfolding.

³Permit and other requirements for the insterstate movement of black stem rust organisms are contained in part 330 of this chapter.

B. sikkimensis	B. thunbergii 'Monry'
B. soulieana 'Claret Cascade'	B. thunbergii 'Painter's Palette'
B. stenophylla	B. thunbergii 'Pink Queen'
B. stenophylla diversifolia	B. thunbergii pluriflora
B. stenophylla gracilis	B. thunbergii 'Royal Burgundy'
B. stenophylla irwini	B. thunbergii 'Royal Cloak'
B. stenophylla nana compacta	B. thunbergii 'Sparkle'
B. taliensis	B. thunbergii 'Thornless'
B. telomaica artisepala	B. thunbergii 'Upright Jewell'
B. thunbergii	B. thunbergii variegata
B. thunbergii 'Antares'	B. thunbergii xanthocarpa
B. thunbergii argenteo marginata	B. thunbergii×'Bailsel' (Golden Car-
B. thunbergii atropurpurea	ousel®)
B. thunbergii atropurpurea erecta	B. thunbergiix'Tara' (Emerald Car-
B. thunbergii atropurpurea erecta	ousel®)
Marshalli	B. triacanthophora
B. thunbergii atropurpurea 'Golden	B. triculosa
Ring'	B. verruculosa
B. thunbergii atropurpurea 'Inter-	B. virgatorum
media'	B. workingensis
B. thunbergii atropurpurea 'Knight	B. xanthoxylon
_ , , ,	
Burgundy' B. thunbergii atropurpurea nana	B.×carminea 'Pirate King' B.×frikartii 'Amstelveen'
B. thunbergii atropurpurea 'Redbird'	
	(3) All plants, seedlings, seeds, fruits,
B. thunbergii atropurpurea 'Rose	and other plant parts capable of propa-
Glow'	gation from the following rust-resist-
B. thunbergii aurea	ant <i>Mahoberberis</i> and <i>Mahonia</i> species,
B. thunbergii 'Aurea Nana'	except <i>Mahonia</i> cuttings for decorative
B. thunbergii 'Bagatelle'	purposes:
B. thunbergii 'Bailgreen' (Jade Car-	(i) Genus <i>Mahoberberis:</i>
ousel TM)	M. aqui-candidula
B. thunbergii 'Bailone'	M. aquifolium 'Smaragd'
B. thunbergii 'Bailone' (Ruby Car-	M. aqui-sargentiae
ousel®)	M. miethkeana
B. thunbergii 'Bailtwo'	M.×'Magic'
B. thunbergii 'Bailtwo' (Burgundy	(ii) Genus <i>Mahonia:</i>
Carousel®)	M. amplectens
B. thunbergii 'Bonanza Gold'	M. aquifolium
B. thunbergii 'Concorde'	M. aquifolium atropurpurea
B. thunbergii 'Crimson Pygmy'	M. aquifolium compacta
B. thunbergii 'Criruzam' Crimson	M. aquifolium compacta 'John Muir'
Ruby TM	M. aquifolium 'Donewell'
B. thunbergii 'Dwarf Jewell'	M. aquifolium 'Kings Ransom'
B. thunbergii erecta	M. aquifolium 'Orangee Flame'
B. thunbergii 'globe'	M. aquifolium 'Undulata'
B. thunbergii 'golden'	M. aquifolium 'Winter Sun'
B. thunbergii 'Golden Pygmy'	M. 'Arthur Menzies'
B. thunbergii 'Green Carpet'	M. bealei
B. thunbergii 'Harlequin'	M. dictyota
B. thunbergii 'Helmond Pillar'	M. fortunei
B. thunbergii 'Kobold'	M. 'Golden Abundance'
B. thunbergii 'Lime Glow'	M. japonica
B. thunbergii 'Lustre Green'	M. japonica×M. lomariifolia 'Charity'
B. thunbergii maximowiczi	M. lomarifolia
B. thunbergii 'Midruzam' Midnight	M. nervosa
$Ruby^{TM}$	M. pinnata
B. thunbergii minor	M. pinnata 'Ken Hartman'
B. thunbergii 'Monlers'	M. piperiana
B. thunbergii 'Monomb'	M. pumila

§301.38-3

M. repens M.×media 'Charity' M.×media 'Winter Sun'

- (4) All plants, seeds, fruits, and other plant parts capable of propagation from rust-susceptible species and varieties of the genera *Berberis, Mahoberberis,* and *Mahonia,* and seedlings from rust-susceptible species and varieties of the genera *Mahoberberis* and *Mahonia,* except *Mahonia* cuttings for decorative purposes.
- (5) Any other product or article not listed in paragraphs (a)(1) through (a)(4) of thissection that an inspector determines presents a risk of spread of black stem rust. The inspectormust notify the person in possession of the product or article that it is subject to the provisions ofthis subpart.
- (b) A person may request that an additional rust-resistant variety be added to paragraph (a)(2) or (a)(3) of this section. The person requesting that a rust-resistant variety be added to paragraph (a)(2) or (a)(3) of this section must provide APHIS with a description of the variety, including a written description and color pictures that can be used by an inspector to clearly identify the variety and distinguish it from other varities.

(Approved by the Office of Management and Budget under control number 0579–0186)

[67 FR 8179, Feb. 22, 2002]

§301.38-3 Protected areas.

(a) The Administrator may designate as a protected area in paragraph (c) of this section any State that has eradicated rust-susceptible plants of the genera Berberis, Mahoberberis, Mahonia under the cooperative Federal-State eradication program. In addition, the State must employ personnel with responsibility for the issuance and withdrawal of certificates in accordance with §301.38-5, and maintain and enforce an inspection program under which every plant nursery within the State is inspected at least once each year to ensure that they are free of rust-susceptible plants. During the requisite nursery inspections, all nursery stock shall be examined to determine that it consists only of rust-resistant varieties of the genera Berberis, Mahoberberis, and Mahonia, and that

the plants are true to type. Plants that do not meet this criteria must be destroyed.

- (b) The Administrator may designate as a protected area any county within a State, rather than the entire State, if areas within the State have eradicated rust-susceptible plants of the genera *Berberis, Mahoberberis,* and *Mahonia* under the cooperative Federal-State program, and;
- (1) The State employs personnel with responsibility for the issuance and withdrawal of certificates in accordance with § 301.38–5;
- (2) The State is enforcing restrictions on the intrastate movement of the regulated articles that are equivalent to those imposed by this subpart on the interstate movement of regulated articles, as determined by the Administrator; and
- (3) The State maintains and enforces an inspection program under which every plant nursery within the county is inspected at least once each year to ensure that plant nurseries within that area are free of rust-susceptible plants of the genera *Berberis, Mahoberberis,* and *Mahonia.* During the requisite nursery inspections, all nursery stock shall be examined to determine that it consists only of rust-resistant varieties of the genera *Berberis, Mahoberberis,* and *Mahonia,* and that the plants are true to type. Plants that do not meet this criteria must be destroyed.
- (c) All seed used to propagate plants of the genera *Berberis, Mahoberberis,* and *Mahonia* in protected areas, and all seed used to propagate plants of the genera *Berberis, Mahoberberis,* and *Mahonia* that are certified as rust-resistant for interstate movement into protected areas, must be produced at properties where a State inspector has verified that no wild or domesticated rust-susceptible plants are growing at or within one-half mile of the property.⁴

⁴Persons performing the inspections must be able to recognize rust-susceptible varieties of *Berberis*, *Mahoberberis*, and *Mahonia*. Inspectors must work side by side, 10 to 20 feet apart, and walk outward away from the property a distance of one-half mile measured from the edge of the property, and observe all plants growing in the half-mile band. The distance between the inspectors